

*Linguistic Mapping*

# The Principles of Calculus I

II

Transformation

II.5

Inverse Functions

*Classroom Exercises*


Copyright © 2024 by Bryan Carrillo and David Weisbart.

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from Bryan Carrillo and David Weisbart.

## Exercise 1


Determining the midpoint  $m$  of a line segment  $L$  with endpoints  $(a, b)$  and  $(c, d)$  comes down to translating an intuitive statement in English to formal mathematics!

To see this, translate this statement into mathematical symbols: Add half the difference between  $(c, d)$  and  $(a, b)$  to move  $(a, b)$  to the midpoint  $m$  of  $L$ .



## Exercise 2


For any line  $L$ , denote by  $\text{Ref}_L$  the reflection across  $L$ . Denote by  $\text{Ref}_1$  rather than  $\text{Ref}_{\text{pow}_1}$  the reflection across  $\text{pow}_1$  to simplify typography. Sketch the line  $\text{pow}_1$  and for any real number  $a$  the points  $(a, 0)$  and  $(0, a)$ . How do you think that  $(a, 0)$  and  $(0, a)$  should reflect across  $\text{pow}_1$ ?



## Exercise 3

For any nonzero real number  $a$ , determine the slope of the line  $L_{\perp}$  that contains  $(a, 0)$  and  $(0, a)$  and determine the midpoint of the line segment with endpoints  $(a, 0)$  and  $(0, a)$ . Use these values to argue that


$$\text{Ref}_1(a, 0) = (0, a) \quad \text{and} \quad \text{Ref}_1(0, a) = (a, 0).$$



## Exercise 4

Determine the midpoint  $m$  of the line segment  $L_{\perp}$  with endpoints  $(a, b)$  and  $(b, a)$  as well as the slope of  $L_{\perp}$ . Conclude that

$$\text{Ref}_1(a, b) = (b, a).$$



## Exercise 5


For any function  $f$  that is a bijection from  $\mathcal{D}(f)$  to  $\mathcal{R}(f)$ ,  $\text{Ref}_1(f)$  is a function from  $\mathcal{R}(f)$  to  $\mathcal{D}(f)$ . Denote by  $f^{-1}$  the function  $\text{Ref}_1(f)$ .

(a) A line  $L$  contains the point  $(1, 3)$  and has slope 5. Given any distinct points  $(a, b)$  and  $(c, d)$  in  $L$ , if  $L$  has slope 5, then

$$\frac{d - b}{c - a} = \boxed{\phantom{0000}}.$$

(b) Compute  $\text{Ref}_1(a, b)$  and  $\text{Ref}_1(c, d)$  to determine the slope of  $L^{-1}$ .

(c) Determine an equation for  $L^{-1}$ .




## Exercise 6

Determining a formula for an inverse function amounts to changing the words one uses to label the components of a function. As an example, take  $f$  to be the function that is given by

$$f(x) = \frac{3x - 1}{x + 2}.$$

- (a) Describe  $f$  using set builder notation and compute  $\text{Ref}_1(f)$ .
- (b) Rewrite  $\text{Ref}_1(f)$  so that the first component is given by an independent variable and the second component depends on the first.








## Exercise 7

The function  $\text{pow}_2$  is not invertible.

- (a) Do you see why  $\text{pow}_2$  is not invertible on  $\mathbb{R}^2$ ? What does  $\text{Ref}_1(\text{pow}_2)$  look like?
- (b) Formulate a precise argument to show that  $\text{pow}_2$  is not invertible.







## Exercise 8


Take  $S$  to be the union of intervals

$$S = (-3, -2] \cup [0, 2) \cup [3, \infty).$$

(a) Do you see why  $\text{pow}_2$  is invertible on  $S$ ? What does  $\text{Ref}_1(\text{pow}_2|_S)$  look like?

(b) Without making a single calculation, present  $(\text{pow}_2|_S)^{-1}$  as a piecewise function.





## Exercise 9

Take  $S$  to be the union of intervals

$$S = (-3, -2] \cup [0, 2) \cup [3, \infty).$$

- Identify a partition  $P$  for  $S$  with as few intervals as possible.
- For each interval  $I$  in  $P$ , determine the range of  $\text{pow}_2 \big|_I$  and a formula for  $\left(\text{pow}_2 \big|_I\right)^{-1}$ .
- Identify a partition  $P'$  for the domain of  $\left(\text{pow}_2 \big|_I\right)^{-1}$  so that  $\left(\text{pow}_2 \big|_I\right)^{-1}$  is given by a single formula on each interval in  $P'$ .
- Present  $\left(\text{pow}_2 \big|_S\right)^{-1}$  as a piecewise defined function with respect to  $P'$ .

